

UNCLASSIFIED

DECLASSIFIED
SIFTED



ILA- 318


~~REDACTED~~ C.3

CIC-14 REPORT COLLECTION
REPRODUCTION
COPY

Classification changed to UNCLASSIFIED
by authority of the U. S. Atomic Energy Commission

Per J.S. Cucchiara, DOC Wash, 9-25-74

By REPORT LIBRARY Bob Martiny, 10-1-74



LOS ALAMOS NATIONAL LABORATORY
3 9338 00350 0781


DECLASSIFIED
SIFTED

UNCLASSIFIED

UNCLASSIFIED

000000
000000

UNCLASSIFIED




REPRODUCTION
OF A SPECIAL COLLECTION
COPY



000000
000000

SECRET

UNCLASSIFIED



UNCLASSIFIED

VERIFIED UNCLASSIFIED

Per JLW 11-26-79
By Linda Kolar

PUBLICLY RELEASABLE

Per M. Jain, FSS-18 Date: 3-27-96
By Kolar, CIC-14 Date: 4-11-96

LA REPORT 318



C.3

June 23, 1945

This document contains 9 pages

APPROXIMATE TREATMENT OF EXPONENTIAL SHOCK AND RAREFACTION WAVES

WORK DONE BY:

R. E. Marshak

REPORT WRITTEN BY:

R. E. Marshak



This document contains information of a classified nature. The disclosure of its contents in any manner to an unauthorized person is prohibited by law.

SECRET


UNCLASSIFIED

SECRET

UNCLASSIFIED


U. S. G. O.
S. I. T. O.
-2-

UNCLASSIFIED



ABSTRACT

It is shown how approximate solutions for the exponential shock and rarefaction waves may be obtained with a minimum of effort. Many such solutions are needed for a discussion of the radiation and mixing phenomena associated with the explosion of the gadget. Figs. 1 to 3 contain some results on the isothermal rarefaction wave.



UNCLASSIFIED
UNCLASSIFIED

S. I. T. O.
U. S. G. O.

UNCLASSIFIED

SECRET
-3-

UNCLASSIFIED

APPROXIMATE TREATMENT OF EXPONENTIAL SHOCK AND RAREFACTION WAVES

The exponential shock and rarefaction waves were treated rigorously by Serber in LA-14 for arbitrary γ (γ is ratio of specific heats at constant pressure and volume). Further in LA-10, Frankel and Davis give the results of a series of numerical integrations for various shock and rarefaction waves. The numerical integrations carried out in LA-10 -- especially for the rarefaction wave -- were quite laborious and for many discussions of the hydrodynamics of the explosion, in particular in connection with radiation and mixing phenomena, it is necessary to get a wide assortment of shock and rarefaction solutions not previously obtained. It is the purpose of this note to discuss briefly how approximate solutions for exponential shock and rarefaction waves may be obtained and to give some results.

We consider first the rarefaction wave (we use Serber's notation (cf. LA-14) throughout); we write:

$$X = X_{-1} e^{\alpha t} \phi(\xi) \tag{1}$$

$$p = p_{-1} e^{2\alpha t} f(\xi) \tag{2}$$

$$\xi = x/X_{-1} e^{\alpha t} \tag{3}$$

where X, x are the Eulerian and Lagrangian positions respectively, p the pressure, t the time and X_{-1}, p_{-1} , and α are constants. The head of the rarefaction wave is taken at $\xi = -1$, and the unit of pressure is chosen so that $f(-1) = 1$. The boundary conditions at the head of the wave are that the displacement is zero, the density normal, and the rarefaction wave moves with the velocity of sound; these conditions imply that

$$\phi(-1) = -1 \tag{4}$$

$$\phi'(-1) = 1 \tag{5}$$

$\frac{p_{-1}}{\rho_{-1} X_{-1}^2 \alpha^2} \gamma_{-1}$ UNCLASSIFIED

UNCLASSIFIED

U. S. G. O.
 OFFICE

UNCLASSIFIED

In (6) ρ_{-1} is the uncompressed density of the core and V_{-1} is the V of the core.

The equations of motion and energy conservation become;

$$\xi^2 \phi'' + \phi - \xi \phi' = - (1/V_{-1}) f' \quad (7)$$

$$(2f - \xi f') \phi' - V_{-1} \xi f \phi'' = 2 \quad (8)$$

In (8) use has also been made of the fact that the pressure is continuous across the head of the rarefaction wave.

If one examines (7) and (8), one finds that there is a sheaf-type singularity at $\xi = -1$, i.e., one coefficient in the power series expansion of say, ϕ_0 is ~~not~~ arbitrary, so that one can get any of an infinite number of solutions depending on the choice of its value. Moreover, since all solutions emerge from the same point $\xi = -1$, it is somewhat difficult to stay on a particular solution if one starts integrating from $\xi = -1$. The more stable method of integration would be to start at $\xi = 0$; but since (7) and (8) constitute a third-order differential equation for ϕ_0 which at most can be reduced to a second-order equation, and since the boundary conditions are all given at $\xi = -1$, this would become too complicated. However, if one examines the solutions in LA-10, one notices that the temperature never changes more than about 5 to 10%. We therefore assume that the temperature is constant throughout the rarefaction wave; this is equivalent to the assumption that $V_{-1} = 1$ and Eqs. (7) and (8) become:

$$\xi^2 \phi'' + \phi - \xi \phi' = - f' \quad (9)$$

$$f \phi' = 1 \quad (10)$$

Inserting (10) into (9) yields the second-order equation:

$$\phi'' (\xi^2 - 1/\phi'^2) + \phi - \xi \phi' = 0 \quad (11)$$

Eq. (11) can be converted into a first-order equation by treating ϕ as the independent variable and $\chi = [\xi \phi' - \phi]$ as the dependent variable; we get:

U. S. G. O.
 OFFICE

UNCLASSIFIED

SECRET
 SECRET
 -5-

$$\frac{dx}{d\phi} = \frac{\gamma(\phi+\chi)}{(\phi+\chi)^2 - 1} \quad (12)$$

Solutions of Eq. (12) are obtained by assuming a value of ϕ at $\xi = 0$ which we denote by ϕ_0 ; this determines χ at $\xi = 0$, namely as $(-\phi_0)$ (since ϕ' is finite at $\xi = 0$). Eq. (12) is then integrated from $\phi = \phi_0$ to $\phi = -1$ and $\phi'(\xi)$ is computed from the relation:

$$\phi'(\xi) = \exp \int_{-1}^{\phi(\xi)} \frac{\chi d\phi}{[(\phi+\chi)^2 - 1]} \quad (13)$$

These integrations go very rapidly and some results are presented in Figs. 1 to 3. Figs. 1 and 2 give ϕ and ϕ' as functions of ξ for various assumed ϕ_0 's. Fig. 3 is a graph of $\phi'(0)$ as a function of ϕ_0 . The solution which is to be used depends on the tamper density; e.g., the solution $\phi_0 = .4741$ fits on to a shock wave with $\gamma = 1$ for equal core and tamper densities. It is interesting to note that the .4741 solution leads to a density at the interface of .45 instead of .44 as in Serber's rigorous treatment.

The exponential shock wave developed in the tamper may be treated approximately in a very simple fashion. Again using Serber's notation (cf. LA-14), we write:

$$x = x_1 e^{\alpha t} \phi(\xi) \quad (14)$$

$$p = p_1 e^{2\alpha t} f(\xi) \quad (15)$$

$$\xi = x/x_1 e^{\alpha t} \quad (16)$$

The point $\xi = 1$ corresponds to the head of the shock wave and there $\phi(1) = 1$, $\phi'(1) = (\gamma-1)/(\gamma+1)$, and $f(1) = 1$. Also, the Hugoniot condition relating the shock velocity to the pressure at the front requires that $p_1/p_0 = [2/(\gamma+1)] \alpha^2 x_1^2$ (p_1 is the tamper density). The pressure and entropy (energy) equations now become;

$$\xi^2 \phi'' + \phi - \xi \phi' = - [2/(\gamma+1)] f' \quad (17)$$

$$f \phi' = \left(\frac{\xi-1}{\gamma+1} \right)^2 \xi^2 \quad (18)$$

SECRET
SECRET
-6-

Since f does not vary much from the front to the back of the shock wave, let us assume that it is constant (and equal to its value at the front) in Eq. (18). Consequently:

$$\phi' = \left(\frac{\gamma-1}{\gamma+1} \right) \xi^{2/\gamma} \quad \text{and} \quad \phi = \frac{\gamma}{(\gamma+2)} \frac{(\gamma-1)}{(\gamma+1)} \xi^{(2+\gamma)/\gamma} + \frac{2(2\gamma+1)}{(\gamma+2)(\gamma+1)} \quad (19)$$

Substituting the above expressions for ϕ and its derivatives into (17), we find for f (using the fact that $f(1)=1$);

$$f = \frac{3\gamma+1}{\gamma+1} - \left(\frac{2\gamma+1}{\gamma+2} \right) \xi - \frac{(\gamma-1)}{(\gamma+1)(\gamma+2)} \xi^{(2+\gamma)/\gamma} \quad (20)$$

Eqs. (19) and (20) constitute an approximate solution for the shock wave. To test the accuracy of the approximation, we evaluate $\phi(0)$ and $f(0)$ for $\gamma = 1.4, 1.67$ and compare with the rigorous values given in LA-10. We get:

$$\begin{aligned} \phi(0) = .931, & \quad f(0) = 2.17 & \quad \text{for} & \quad \gamma = 1.4 \\ \phi(0) = .886, & \quad f(0) = 2.25 & \quad \text{for} & \quad \gamma = 1.67 \end{aligned}$$

The rigorous values are:

$$\begin{aligned} \phi(0) = .944, & \quad f(0) = 2.20 & \quad \text{for} & \quad \gamma = 1.4 \\ \phi(0) = .907, & \quad f(0) = 2.31 & \quad \text{for} & \quad \gamma = 1.67 \end{aligned}$$

The agreement is quite good and moreover it is relatively easy to obtain increasingly better approximations. For example, the next approximation to the specific volume (i.e., ϕ') -- found by substituting (20) into (18) -- yields very accurate results.

SECRET
SECRET
APPROVED FOR PUBLIC RELEASE




FIGURE 1

UNCLASSIFIED


UNCLASSIFIED

APPROVED FOR PUBLIC RELEASE

APPROVED FOR PUBLIC RELEASE

-ξ-

FIGURE 2



APPROVED FOR PUBLIC RELEASE

APPROVED FOR PUBLIC RELEASE





FIGURE 3


UNCLASSIFIED



APPROVED FOR PUBLIC RELEASE

APPROVED FOR PUBLIC RELEASE

UNCLASSIFIED



DECLAS
SIFIED

UNCLASSIFIED

UNCLASSIFIED

DOCUMENT ROOM

REC. FROM OS

DATE 6/26/45

REC. NO. REC. ✓

UNCLASSIFIED

UNCLASSIFIED

DECLAS

SIFIED