


LA -- 1258
Copy 1 of 115
Series..A

PHOTO MATL. LAB. LIBS.



3 9338 00419 7439

UNCLASSIFIED
UNCLASSIFIED

0377

LOS ALAMOS SCIENTIFIC LABORATORY


of

THE UNIVERSITY OF CALIFORNIA

May 26, 1951

LA-158

This document consists of 10 pages



PUBLICLY RELEASABLE
LANL Classification Group
Atgam 11/16/95

THE NEUTRON INDUCED FISSION CROSS SECTION OF U²³⁶ AS A
FUNCTION OF ENERGY

Classification changed to UNCLASSIFIED
by authority of the U. S. Atomic Energy Commission.
Per *Charles F. Speed 10/1/60*
W. N. Nyer 5-16-60
By REPORT LIBRARY

Work done by:

Warren Nyer

Report written by:

Warren Nyer


LOS ALAMOS NATIONAL LABORATORY



3 9338 004 19 7439

PHYSICS - FISSION

0377



UNCLASSIFIED

UNCLASSIFIED



PHYSICS - FISSION

JUN 19 1951


LA-1258

Los Alamos Document Room
J R. Oppenheimer

1-20
21

STANDARD DISTRIBUTION

Argonne National Laboratory	22-33
Armed Forces Special Weapons Project	34
Atomic Energy Commission, Washington	35-40
Brookhaven National Laboratory	41-44
Bureau of Ships	45
Carbide & Carbon Chemicals Company (K-25 Plant)	46-47
Carbide & Carbon Chemicals Company (Y-12 Area)	48-49
Chicago Patent Group	50
Chief of Naval Research	51
Columbia University (Havens)	52
duPont Company	53-56
General Electric Company, Richland	57-60
Hanford Operations Office	61
Idaho Operations Office	62-63
Iowa State College	64
Knolls Atomic Power Laboratory	65-68
Mound Laboratory	69-70
National Advisory Committee for Aeronautics	71
Naval Research Laboratory	72
New York Operations Office	73-74
North American Aviation, Inc.	75-77
Oak Ridge National Laboratory (X-10)	78-85
Patent Branch, Washington	86
Savannah River Operations Office	87
UCLA Medical Research Laboratory (Warren)	88
USAF - Headquarters	89
U. S. Naval Radiological Defense Laboratory	90
University of California Radiation Laboratory	91-93
Westinghouse Electric Corporation	94-97
Wright-Patterson Air Force Base	98-100
Technical Information Service, Oak Ridge	101-115



UNCLASSIFIED

~~CONFIDENTIAL~~

- 3

ABSTRACT

The neutron-induced fission cross section for the isotope U²³⁶ has been measured relative to the fission cross section for the isotope U²³⁵ as a function of neutron energy. The threshold of the reaction is in the neighborhood of 690 kev. The cross section rises to 0.69 barns at 1.3 Mev. At 2.55 Mev it is 0.53 barns, and at 14.1 Mev it is 1.65 barns.

UNCLASSIFIED

03115

03115

- 4 -

The neutron-induced fission cross section of 261 has been

-
1. The isotopes are identified throughout by a code using the combination of the terminal number of the atomic number and the terminal number of the atomic weight, thus thorium 232 is 02 and uranium 234 is 25.
-

measured relative to the neutron-induced fission cross section of 25 as a function of energy using a parallel-plate comparison chamber² in which a 26

2. LAMS 938
-

foil and a 25 foil were placed back-to-back. The chamber was filled with argon to a pressure of 70 cm of mercury and carbon dioxide to a pressure of 5 cm of mercury. Electron collection was employed. Model 502 fast amplifiers were used to minimize "pile-up" of pulses arising from α particles.

The measurements consisted of determining the ratio R_E of the number of fissions in the 26 foil³ to the number of fissions in the 25 foil³ when the chamber was placed in a flux of neutrons of known energy E . Except

3. The foils were prepared by John Povelites of Group CMR-4
-

in negligibly few cases, every fission resulted in an ionization pulse in the chamber. A measurement of the ratio R_{th} at thermal energies in the standard graphite pile served to "weigh" the 26 foil, since the isotopic composition of the foils and the mass of the 25 foil is known and it can be assumed that only the 25 undergoes thermal fission. This procedure eliminates some of the uncertainties involved in the preparation and use of a rare isotope foil. The weight of the 26 foil deduced by the method just indicated agreed within 1.5% with the mass values obtained by balance weighing and by α -counting.

UNCLASSIFIED

• • • • •

• • • • •
 • • • • •
 • • • • •

Van de Graaff acceleration in W Building using the $T(p,n)He^3$ reaction.

The spread in energy of the neutrons from this source was calculated from the thickness of the tritium gas target and the spread in angle between the neutron and the proton beam, which was from 0° to 20° . The energy assigned to each measurement was a value such that half the neutrons had energy greater and half the neutrons energy less than this value. The error in the neutron energy due to uncertainties in the energy loss of protons in aluminum and tritium is about ± 20 kev. No estimate of the errors in the cross section due to the presence of low energy epi-cadmium neutrons was made for any measurement except the 14.1 Mev case. It is believed that this error is small. In all measurements the chamber was surrounded with cadmium to eliminate thermal neutrons. The Van de Graaff accelerator was operated by the P-3 accelerator crew.

The Z Building Cockcroft-Walton accelerator was used to produce the 2.55 and 14.1 Mev neutrons by means of the $D(D,n)He^3$ and $T(D,n)He^4$ reactions, respectively. The measurements were made at 90° to the beam. The energy spread due to angle spread was 160 kev for the 2.55 Mev case and 190 kev for the 14.1 Mev case. A correction of about 3% was applied for the effect of low energy epi-cadmium neutrons in the 14.1 Mev measurement. The magnitude of this correction had been determined by a 25 spiral chamber measurement.² Where necessary, corrections were made for the difference in flux at the two foils. These corrections were of the order of a few percent.

The cross sections obtained and the statistical errors are given in Table 1. The 25 and 28 cross sections were taken from the smooth curves

• • • • •
 • • • • •
 • • • • •
 • • • • •
 • • • • •

• • • • •
- 7 -
• • • • •
• • • • •

in LA-520 for energies below 14 Mev. The 14 Mev values of these cross sections were taken from LA-719 and LAMS-938. The plus and minus values on the neutron energies listed in Table I indicate the spread in neutron energy.

• • • • •
• • • • •
• • • • •
• • • • •





TABLE I


FISSION CROSS SECTION OF U²³⁶ AS A FUNCTION OF ENERGY


Neutron Energy	No of 26 Fission Counts	Corrected Ratio of 26 Fissions to 25 Fissions	$\frac{\sigma_f(26)}{\sigma_f(25)}$	$\sigma_f(26)$ (barns)
Thermal	14,178	0.203		
690 ± 34 kev	1,040	0.207	0.03 ± 0.6	0.04 ± .08
770 * 35 - 40 kev	1,088	0.221	0.14 ± .06	0.19 ± .08 ←
1000 * 40 - 50 kev	1,868	0.238	0.27 ± .05	0.36 ± .07
1320 * 33 - 52	1,014	0.271	0.52 ± .07	0.69 ± .10
2.55 ± .08 Mev	1,155	0.259	0.40 ± .04	0.53 ± .06
14.1 ± .1 Mev	4,684	0.298	0.71 ± .03	1.65 ± .10

*65 a la. Sample ORNL
Sept '52*

*47 = σ ~ 8 mega used
55*







SECRET

DOCUMENT ROOM

REC. BY SA

DATE 6-16-51

REC. NO. REC. _____

SECRET